

Has the QCD Critical Point been Signaled by Observations at RHIC ?

Roy A. Lacey, N. N. Ajitanand, J. M. Alexander, P. Chung, W.G. Holzmann, M. Issah, and A. Taranenko
Department of Chemistry, State University of New York at Stony Brook, Stony Brook, NY 11794-3400, USA

P. Danielewicz¹ and Horst Stöcker²

¹*National Superconducting Cyclotron Laboratory and Department of Physics and Astronomy,
 Michigan State University, East Lansing, MI 48824-1321, USA*

²*Institut für Theoretische Physik, Johann Wolfgang Goethe-Universität D60438 Frankfurt am Main, Germany
 (Dated: February 5, 2008)*

The shear viscosity to entropy ratio (η/s) is estimated for the hot and dense QCD matter created in Au+Au collisions at RHIC ($\sqrt{s_{NN}} = 200$ GeV). A very low value is found $\eta/s \sim 0.1$, which is close to the conjectured lower bound ($1/4\pi$). It is argued that such a low value is indicative of thermodynamic trajectories for the decaying matter which lie close to the QCD critical end point.

PACS numbers: PACS 25.75.Ld

A primary objective for studying ultra-relativistic heavy ion collisions is to map out the accessible regions of the Quantum Chromo Dynamics (QCD) phase diagram. The existence of a critical point – an end point of the first order chiral transition in this diagram – has attracted considerable attention since it was first proposed [1, 2]. Several theoretical techniques have been exploited to locate this critical end point (CEP) in the plane of temperature vs baryon chemical potential (T, μ_B) i.e. the QCD phase diagram [3, 4, 5, 6, 7]. The properties of the CEP have also been sought via model calculations and the universality hypothesis [8, 9, 10]. Nonetheless, experimental verification of the CEP constitutes a major current scientific challenge.

The study of heavy ion collisions to search for the CEP was proposed several years ago [11]. Recently, a resurgence of experimental interest has centered around the prospects for successful experimental searches at Brookhaven's Relativistic Heavy Ion Collider (RHIC) [12, 13] and CERN's Super Proton Synchrotron (SPS) [14, 15]. Indeed, a recent theoretical investigation involving two flavors of light dynamical quarks [16] gives $T_c/T_{co} \sim 0.95$ and $\mu_B^c/T_{co} \sim 1.1$, where $T_{co} \sim 170$ MeV is the temperature for a cross-over to the quark gluon plasma (QGP) at zero baryon chemical potential and T_c and μ_B^c are the temperature and the baryon chemical potential (respectively) at the critical end point. It is widely believed that these values for T_c and μ_B^c place the CEP in the range of observability for future energy scans at both RHIC and the SPS [13].

In recent hydrodynamical calculations [17, 18], an interesting aspect of the CEP has been reported which could have important consequences for its detection. That is, it acts as an attractor for thermodynamic trajectories of evolving hot and dense QCD matter [11]. This aspect of the CEP is illustrated in Fig. 1 where isentropic trajectories in the (T, μ_B) plane are shown for a broad range of values for the entropy per baryon s/n_B . Trajectories for the range $s/n_B \sim 100 - 50$, are clearly focused




FIG. 1: (Color online) Results from a hydrodynamical calculation by Nonaka et al which assumes an EOS with a CEP [17]. Isentropic trajectories are shown for the case in which the CEP is located at $(T, \mu_B) = (154.7 \text{ MeV}, 367.8 \text{ MeV})$. The value of s/n_B for each of these trajectories (left to right) is 100.0, 66.6, 50.0, 40, 33.3, 28.6 and 25.

toward the CEP. Such a focusing is not observed for isentropic trajectories in similar model calculations which do not take explicit account of the CEP in the equation of state (EOS). Instead, trajectories are merely shifted on the phase coexistence line [17]. Given this aspect of the CEP, it is worthwhile to investigate whether or not the decay dynamics of the high energy density QCD matter created in heavy ion RHIC collisions show any signals for the CEP.

Here, we argue that estimates for the ratio of viscosity to entropy density (η/s) made from RHIC measurements [19, 20, 21], indicate decay trajectories which are close to the CEP. This implies that the prospects for constraining the location of the CEP in the (T, μ_B) plane are very good. Possibly, such constraints can be best achieved via a study of excitation functions with relatively large steps in collision energy prior to focusing on a series of fine steps.




FIG. 2: (Color online) $v_2/\varepsilon n_q$ vs KE_T/n_q for several identified particle species obtained in minimum bias Au+Au collisions. The data is taken from Refs. [32, 33] and the references therein. The dashed-dotted line represents a fit to the data.

There is strong experimental evidence for the creation of locally equilibrated nuclear matter at unprecedented energy densities, in heavy ion collisions at RHIC [22, 23, 24, 25, 26, 27, 28, 29]. Jet suppression studies indicate that the constituents of this matter interact with unexpected strength and this matter is almost opaque to high energy partons [30, 31].

Elliptic flow measurements [32, 33] validate the predictions of perfect fluid hydrodynamics for the scaling of the elliptic flow coefficient v_2 with eccentricity ε , system size and transverse kinetic energy KE_T [34, 35, 36]; they also indicate the predictions of valence quark number (n_q) scaling [37, 38, 39], suggesting that quark-like degrees of freedom are pertinent when elliptic flow develops [32, 33]. The result of such scaling is illustrated in Fig. 2 where we plot $v_2/n_q\varepsilon$ vs KE_T/n_q ; it shows that the relatively “complicated” dependence of v_2 on centrality, transverse momentum, particle type and quark number can be scaled to a single function [32, 33].

The expected suppression of flow due to the shear viscosity, as predicted by weak-coupling transport calculations is also not observed. In fact, only a rather small viscosity, $\eta/s \lesssim 0.1$, could be accommodated by the data in early estimates [19]. This value of η/s , which is much lower than that obtained from weak coupling QCD [40] or hadronic computations [41], has been interpreted as evidence that the quark gluon plasma (QGP) created in the early phase of RHIC collisions is more strongly coupled than expected [28, 42]. An alternative interpretation is

that the low value for η/s could result from an anomalous viscosity η_A , arising from turbulent color magnetic and electric fields dynamically generated in the expanding quark-gluon plasma [43]. That is, $\eta^{-1} = \eta_A^{-1} + \eta_C^{-1}$ and η_A dominates over the collisional viscosity η_C .

The ratio η/s cannot be arbitrarily small because quantum mechanics limits the size of cross sections via unitarity. A conjectured lower bound for η/s is $1/4\pi$, reached in the strong coupling limit of certain gauge theories [44]. It has even been speculated that this lower bound holds for all substances [44]. The temperature dependence of η/s also provides invaluable insights. This is illustrated in Fig. 3 where we have plotted η/s vs $(T - T_c)/T_c$ for molecular, atomic and nuclear matter. The data for He, N₂ and H₂O were obtained for their respective critical pressure. They are taken from Ref. [45] and the references therein. The calculated results shown for the meson-gas (for $T < T_c$) are obtained from chiral perturbation theory with free cross sections [46]. Those for the QGP (ie. for $T > T_c$) are from lattice QCD simulations [47]. The value $T_c \sim 170$ MeV is taken from lattice QCD calculations [48].

Figure 3 illustrates the observation that for atomic and molecular substances, the ratio η/s exhibits a minimum of comparable depth for isobars passing in the vicinity of the liquid-gas critical point [44, 45, 46]. When an isobar passes through the critical point (as shown in Fig. 3), the minimum forms a cusp at T_c ; when it passes below the critical point, the minimum is found at a temperature below T_c (liquid side) but is accompanied by a discontinuous change across the phase transition. For an isobar passing above the critical point, a less pronounced minimum is found at a value slightly above T_c . The value η/s is smallest in the vicinity of T_c because this corresponds to the most difficult condition for the transport of momentum [45].

Given the observations for atomic and molecular substances, one expects a broad range of trajectories in the (T, n_B) or (T, μ_B) plane for nuclear matter, to show η/s minima with a possible cusp at the critical point. The exact location of this point (in the (T, μ_B) plane) is of course not known, and only coarse estimates of where it might lie are available. The open triangles in Fig. 3 show calculated values for η/s along the $\mu_B = 0$, $n_B = 0$ trajectory. For $T < T_c$ the η/s values for the meson-gas show an increase for decreasing values of T . For T greater than T_c , the lattice results [47] indicate an increase of η/s with T , albeit with large error bars. These trends suggest a minimum for η/s .

The open hexagons in Fig. 3 show estimates for η/s constrained by experimental data; they are for a trajectory which starts along $n_B = n_0$. The η values are extracted from momentum transport data obtained in intermediate-energy (IE) collisions [49, 50, 51]; values for s were obtained from composite yields measured in the same energy and mass region. The filled hexagon in




FIG. 3: (Color online) η/s vs $(T - T_c)/T_c$ for several substances as indicated. The calculated values for the meson-gas have an associated error of $\sim 50\%$ [46]. The lattice QCD value $T_c = 170$ MeV [48] is assumed for nuclear matter. The lines are drawn to guide the eye.

Fig. 3 shows a recent estimate for η/s from high precision elliptic flow measurements in Au+Au collisions ($\sqrt{s_{NN}} = 200$ GeV) at RHIC [21]:

$$\frac{\eta}{s} \sim T \lambda_f c_s,$$

where T is the temperature, λ_f is the mean free path and c_s is the sound speed in the matter.

The temperature $T = 160$ MeV was employed in a blast wave model generalized to include viscous corrections [19]. We estimate the value $T = 165 \pm 3$ MeV via a hydrodynamically inspired fit to the universally scaled v_2 data shown in Fig. 2. The functional form used for the fit is $I_1(w)/I_0(w)$, where $w = KE_T/2T$ and $I_1(w)$ and $I_0(w)$ are Bessel functions [35]. For c_s , we use the recent observation that v_2 scales with eccentricity and across system size to constrain the estimate $c_s = 0.35 \pm 0.05$ [32, 33, 52]. The mean free path estimate $\lambda_f = 0.3 \pm 0.03 fm$ [53], is obtained from an on-shell transport model simulation of the gluon evolution (in space and time) in Au+Au collisions at 200 GeV [54]. This parton cascade model includes pQCD $2 \leftrightarrow 2$ and $2 \leftrightarrow 3$ scatterings. These values for λ_f , T and c_s lead to the estimate $\eta/s \sim 0.09 \pm 0.015$ with an estimated systematic error of $+0.1$ primarily due to the uncertainty in λ_f . This value for η/s is in good agreement with the experimentally based estimates of

Teaney and Gavin [19, 20] and the theoretical estimates of Gyulassy and Shuryak [42, 55]. However, it is in stark contrast to the predictions of perturbative QCD [19, 40].

The experimental and calculated η/s values for nuclear matter shown in Fig. 3, give strong indication for a minimum in the vicinity of T_c . This minimum, achieved in RHIC collisions, is rather close to the conjectured lower bound of $\eta/s = 1/4\pi$. We therefore conclude that it is compatible with the minimum expected if the hot and dense QCD matter produced in RHIC collisions follow decay trajectories which are close to the CEP. As discussed earlier, such trajectories could be readily followed if the CEP acts as an attractor for thermodynamic trajectories for the decaying matter [17, 18]. Here, it is noteworthy that the system has to spend a considerable portion of its dynamics in the region of low η/s for that low value to have a significant impact on the reaction dynamics. If the system gets momentarily, to a low η/s and the rest of the dynamics takes place at high η/s , the momentary low value will have little impact. An optimal situation would be if the system stays at low η/s and then very quickly freezes out thereafter. That is, significant collective motion develops prior to the systems evolution toward higher η/s values.

In summary we have used the comprehensive overall scaling of elliptic flow data to constrain a rudimentary experimental estimate for η/s in Au+Au collisions at RHIC [56]. A value close to the conjectured lower bound of $\eta/s = 1/4\pi$ is observed. We argue that such a low value for η/s is suggestive of decay trajectories which lie close to the QCD critical end point, because it acts as an attractor for constant entropy phase space trajectories of evolving hot and dense QCD matter. This suggests that the prospects for a successful search for the first order QCD phase transition and the possibility for establishing reliable constraints (in the (T, μ_B) plane) for its associated critical end point are quite good. An initial measurement of excitation functions with coarse-grained collision energy steps may give sufficient information, or at least, give a better perspective on where to focus attention on fine steps.

This work was supported by the US DOE under contract DE-FG02-87ER40331.A008 and by the U.S. National Science Foundation under Grant No. PHY-0555893.

-
- [1] M. Asakawa and K. Yazaki, Nucl. Phys. **A504**, 668 (1989).
 - [2] A. Barducci, R. Casalbuoni, S. De Curtis, R. Gatto, and G. Pettini, Phys. Lett. **B231**, 463 (1989).
 - [3] Z. Fodor and S. D. Katz, JHEP **03**, 014 (2002), hep-lat/0106002.
 - [4] P. de Forcrand and O. Philipsen, Nucl. Phys. **B673**, 170 (2003), hep-lat/0307020.

- [5] C. R. Allton et al., Phys. Rev. **D71**, 054508 (2005), hep-lat/0501030.
- [6] R. V. Gavai and S. Gupta, Phys. Rev. **D71**, 114014 (2005), hep-lat/0412035.
- [7] O. Philipsen, Proceedings of Science **LAT2005**, 016 (2006), hep-lat/0510077.
- [8] J. Berges and K. Rajagopal, Nucl. Phys. **B538**, 215 (1999), hep-ph/9804233.
- [9] M. A. Halasz, A. D. Jackson, R. E. Shrock, M. A. Stephanov, and J. J. M. Verbaarschot, Phys. Rev. **D58**, 096007 (1998), hep-ph/9804290.
- [10] M. A. Stephanov, Prog. Theor. Phys. Suppl. **153**, 139 (2004), hep-ph/0402115.
- [11] M. A. Stephanov, K. Rajagopal, and E. V. Shuryak, Phys. Rev. Lett. **81**, 4816 (1998), hep-ph/9806219.
- [12] RHIC-Workshop (2006), <https://www.bnl.gov/riken/QCDRhic/>.
- [13] G. S. F. Stephens (2006), nucl-ex/0607030.
- [14] M. Gazdzicki and M. I. Gorenstein, Acta Phys. Polon. **B30**, 2705 (1999), hep-ph/9803462.
- [15] M. Gazdzicki (2005), nucl-ex/0512034.
- [16] R. Gavai and S. Gupta, Proceedings of Science **LAT2005**, 160 (2006), hep-lat/0509151.
- [17] C. Nonaka and M. Asakawa, Phys. Rev. **C71**, 044904 (2005), nucl-th/0410078.
- [18] B. Kampfer, M. Bluhm, R. Schulze, D. Seipt, and U. Heinz (2005), hep-ph/0509146.
- [19] D. Teaney, Phys. Rev. **C68**, 034913 (2003), nucl-th/0301099.
- [20] S. Gavin and M. Abdel-Aziz (2006), nucl-th/0606061.
- [21] R. A. Lacey (2006), nucl-ex/0608046.
- [22] K. Adcox et al., Nucl. Phys. **A757**, 184 (2005), nucl-ex/0410003.
- [23] J. Adams et al., Nucl. Phys. **A757**, 102 (2005), nucl-ex/0501009.
- [24] B. B. Back et al., Nucl. Phys. **A757**, 28 (2005), nucl-ex/0410022.
- [25] I. Arsene et al., Nucl. Phys. **A757**, 1 (2005), nucl-ex/0410020.
- [26] M. Gyulassy et al., Nucl. Phys. **A750**, 30 (2005), nucl-th/0405013.
- [27] B. Müller (2004), nucl-th/0404015.
- [28] E. V. Shuryak, Nucl. Phys. **A750**, 64 (2005), hep-ph/0405066.
- [29] U. Heinz et al., Nucl. Phys. **A702**, 269 (2002).
- [30] C. Adler et al. (STAR), Phys. Rev. Lett. **90**, 082302 (2003), nucl-ex/0210033.
- [31] S. S. Adler et al. (PHENIX), Phys. Rev. Lett. **97**, 052321 (2006), nucl-ex/0507004.
- [32] M. Issah and A. Taranenko (PHENIX) (2006), nucl-ex/0604011.
- [33] A. Adare (PHENIX) (2006), nucl-ex/0608033.
- [34] M. Csanád et al. (2005), nucl-th/0512078.
- [35] M. Csanád et al. (2006), nucl-th/0605044.
- [36] R. S. Bhalerao et al., Phys. Lett. **B627**, 49 (2005).
- [37] S. A. Voloshin, Nucl. Phys. **A715**, 379 (2003), nucl-ex/0210014.
- [38] R. J. Fries et al., Phys. Rev. **C68**, 044902 (2003).
- [39] V. Greco, C. M. Ko, and P. Levai, Phys. Rev. **C68**, 034904 (2003), nucl-th/0305024.
- [40] P. Arnold, G. D. Moore, and L. G. Yaffe, JHEP **11**, 001 (2000).
- [41] A. Muronga, Phys. Rev. **C69**, 044901 (2004), nucl-th/0309056.
- [42] T. Hirano and M. Gyulassy, Nucl. Phys. **A769**, 71 (2006), nucl-th/0506049.
- [43] M. Asakawa, S. Bass, and B. Müller (2006), hep-ph/0603092.
- [44] P. Kovtun, D. T. Son, and A. O. Starinets, Phys. Rev. Lett. **94**, 111601 (2005), hep-th/0405231.
- [45] L. P. Csernai, J. I. Kapusta, and L. D. McLerran, Phys. Rev. Lett. **97**, 152303 (2006), nucl-th/0604032.
- [46] J.-W. Chen and E. Nakano (2006), hep-ph/0604138.
- [47] A. Nakamura and S. Sakai, Phys. Rev. Lett. **94**, 072305 (2005), hep-lat/0406009.
- [48] F. Karsch, E. Laermann, and A. Peikert, Nucl. Phys. **B605**, 579 (2001), hep-lat/0012023.
- [49] P. Danielewicz, Acta. Phys. Pol. **B33**, 45 (2002).
- [50] W. Schmidt et al., Phys. Rev. **C47**, 2782 (1993).
- [51] P. Danielewicz and B. Barker, In preparation (2006).
- [52] R. S. Bhalerao and J.-Y. Ollitrault, Phys. Lett. **B641**, 260 (2006), nucl-th/0607009.
- [53] This estimate is consistent with observed universal scaling of v_2 for D mesons presented in Ref. [57] (????).
- [54] Z. Xu and C. Greiner, Phys. Rev. **C71**, 064901 (2005), hep-ph/0406278.
- [55] B. A. Gelman, E. V. Shuryak, and I. Zahed (2006), nucl-th/0601029.
- [56] Further constraints leading to an improved estimate could come from future experimental studies of the scaling violations to perfect fluid flow predictions [45]. (????).
- [57] R. A. Lacey and A. Taranenko (2006), nucl-ex/0610029.